

Recovering of curves with involution by extended Prym data

Vassil Kanev*

Introduction

The classical Torelli theorem states that every smooth, projective algebraic curve X is determined uniquely up to isomorphism by its principally polarized Jacobian $(J(X), \Theta)$. In this paper we consider curves $\tilde{C}$ with involution without fixed points $\sigma : \tilde{C} \longrightarrow \tilde{C}$. We let $C = \tilde{C}/\sigma$ and denote by $\pi : \tilde{C} \longrightarrow C$ the factor map. One associates the principally polarized Prym variety (P, Ξ) where $P = P(\tilde{C}, \sigma) = (1 - \sigma)J(\tilde{C})$ and $\Theta|_P$ is algebraically equivalent to 2Ξ . The natural question, whether the Prym variety determines uniquely the pair $(\tilde{C}, \sigma)$ up to isomorphism, has negative answer in general if the genus g of C is ≤ 6 as well as in every genus ≥ 7 for some special loci of curves, e.g for hyperelliptic C . The problem of spotting the pairs $(\tilde{C}, \sigma)$ which are not determined uniquely by the Prym variety is still open. Some partial results have been obtained in [F-S],[K],[Do1],[Do2],[De1],[De2],[N],[V].

In this paper we propose an extension of the Prym data (P, Ξ) and prove that it determines uniquely up to isomorphism any pair $(\tilde{C}, \sigma)$ for $g \geq 2$. Our extension originates from the following observation. Consider the case of $\tilde{C}$ of genus 1. Here the involution is a translation t_μ by some point μ of order 2 in $J(\tilde{C})$. Notice that $\{0, \mu\} = \text{Ker}(Nm_\pi : J(\tilde{C}) \longrightarrow J(C))$. The Prym variety is equal to 0. There is a classically known data which determines uniquely the pair $(\tilde{C}, t_\mu)$ up to isomorphism (see e.g. [M3], [C]). Namely, one can always represent $J(\tilde{C})$ as $\mathbf{C}/\mathbf{Z}\tau + \mathbf{Z}$, where τ belongs to the upper-half plane $\mathcal{H}$, so that

$$\mu = \frac{1}{2}\tau + \frac{1}{2}(\text{mod } \mathbf{Z}\tau + \mathbf{Z}).$$

*Supported in part by the Bulgarian foundation "Scientific research" and by NSF under the US-Bulgarian project "Algebra and algebraic geometry".

So, the moduli space of pairs $(\tilde{C}, \sigma)$ is isomorphic to $\Gamma_{1,2} \backslash \mathcal{H}$ where $\Gamma_{1,2} \subset PGL(2, \mathbf{Z})$ is the subgroup

$$\Gamma_{1,2} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid ad - bc = 1, ab \equiv 0 \pmod{2}, cd \equiv 0 \pmod{2} \right\}$$

One considers the three even theta functions with characteristics : $\theta_{00}(z, \tau), \theta_{01}(z, \tau)$ and $\theta_{10}(z, \tau)$. Just one of them vanishes on μ , namely $\theta_{00}(z, \tau)$. Now, by the transformation law for theta functions one checks that for

$$\lambda(\tau) = -\theta_{01}(0, \tau)^4 / \theta_{10}(0, \tau)^4 \quad (1)$$

the set $\{\lambda(\tau), 1/\lambda(\tau)\}$, or equivalently the function $k(\tau) = \lambda(\tau) + 1/\lambda(\tau)$, remain invariant with respect to the action of $\Gamma_{1,2}$. Moreover the map

$$k : \Gamma_{1,2} \backslash \mathcal{H} \longrightarrow \mathbf{C}^* - \{0, 2\}$$

is an analytic isomorphism (see Section (1)).

Generalizing to genus greater than 1, first we have that $Ker(Nm_\pi : J(\tilde{C}) \longrightarrow J(C)) = P \cup P_-$ where P_- is a translation of P by a point of order 2. One considers the symmetric theta divisors Θ of $J(\tilde{C})$ which have the property that for every $\rho \in P_2$ either $\rho \notin \Theta$ or $\rho \in \Theta$ and $mult_\rho \Theta$ is even. It turns out that there are three P_2 -orbits of symmetric theta divisors with this property. The divisors of one of the orbits contain P_- ; these are exactly the theta divisors which appear in Wirtinger's theorem [F],[M2] and satisfy $\Theta.P = 2\Xi$ for symmetric theta divisors Ξ of P . None of the divisors of the other two orbits contains P_- . Restricting the latter to P we obtain two P_2 -orbits of divisors in the linear system $|2\Xi|$ which we denote by O_1, O_2 .

The extended Prym data consists of (P, Ξ) together with the two P_2 -orbits $O_1, O_2 \subset |2\Xi|$. Our result is that for $g \geq 2$ it determines uniquely up to isomorphism any pair $(\tilde{C}, \sigma)$. The proof is analogous to Andreotti's proof of Torelli's theorem and uses the Gauss map for the divisors of O_i . Special treatment is required if C is hyperelliptic, or bi-elliptic, or $g=3$. In fact the bi-elliptic case has been already considered earlier by Naranjo in [N] who proved that for $g \geq 10$ the pair $(\tilde{C}, \sigma)$ can be recovered by the ordinary Prym data (P, Ξ) . His arguments however do not work for $g = 4$ or 5 . If η is the point of order 2 in $Pic^0(C)$ which

determines the covering $\pi : \tilde{C} \longrightarrow C$ then the divisors of O_i are equal to translations of connected components of the set

$$Z = \{L \in Pic^{2g-2}(\tilde{C}) \mid Nm_\pi(L) \simeq K_C \otimes \eta, h^0(\tilde{C}, L) \geq 1\}$$

It is interesting that line bundles L of this type appear also in the study of rank 2 vector bundles on C with canonical determinant [B] as well as in representing $(\tilde{C}, \sigma)$ as the spectral curves associated to $sp(2n)$ -matrices with parameter.

Acknowledgement. Part of this work was done while the author was a visitor in the University of Michigan and the University of Utah. The hospitality of these institutions is gratefully acknowledged.

Contents

0. Notation and preliminaries. 1. Double unramified coverings of elliptic curves. 2. Extended Prym data. 3. The semicanonical map and the Gauss map. 4. The hyperelliptic case, $g \geq 2$. 5. The bi-elliptic case, $g \geq 4$. 6. The case $g = 3$.

0 Notation and preliminaries

We denote by $\equiv$ the linear equivalence of divisors. Let X be an algebraic, smooth, irreducible curve. We denote by $J_d(X)$ the divisor classes of degree d modulo linear equivalence and by $Pic^d(X)$ the isomorphism classes of invertible sheafs of degree d on X . Abusing the notation we shall write by the same letter an element in $J_d(X)$ and the corresponding element in $Pic^d(X)$. If D is a divisor of degree d and $\xi = cl(D)$ its class in $J_d(X)$ we write

$$h^0(C, D) = h^0(C, \xi) = \dim |\xi| + 1$$

If L is an invertible sheaf of X , then $|L|$ is the linear system of divisors of sections of $H^0(X, L)$. If $|L|$ is without base points we denote by $\varphi_L = \varphi_{|L|}$ the map $X \longrightarrow |L|^*$ defined by $\varphi_L(x) = |L(-x)| + x$.

Let A be a principally polarized abelian variety of dimension g isomorphic to $\mathbf{C}^g/\Lambda_\tau$, where $\Lambda_\tau = \mathbf{Z}^g\tau + \mathbf{Z}^g$ with $\tau \in \mathcal{H}_g$, where $\mathcal{H}_g$ is the Siegel upper-half space. Any point $e \in \mathcal{C}^g$

has two characteristics $\epsilon, \delta \in \mathbf{R}^g$ such that $e = \epsilon\tau + \delta$. We shall write sometimes $e = \begin{bmatrix} \epsilon \\ \delta \end{bmatrix}$. Let $x \in A$ and let $x = (x'\tau + x'')(mod \Lambda_\tau)$ with $x', x'' \in \mathbf{R}^g$. If no confusion arises we shall refer to x', x'' as the characteristics of x , keeping in mind that x', x'' are determined modulo $\mathbf{Z}^g$. We shall denote by A_2 the points of order 2 in A . Let $\lambda = \lambda'\tau + \lambda'', \mu = \mu'\tau + \mu'' \in A_2$. The Weyl pairing $e_2 : A_2 \times A_2 \longrightarrow \mathbf{Z}$ is defined by

$$e_2(\lambda, \mu) = 4(\lambda^t \mu'' - \mu^t \lambda''(mod 2))$$

Let Θ be a symmetric theta divisor. One defines a quadratic form $q_\Theta : A_2 \longrightarrow \mathbf{Z}_2$ associated with Θ by

$$q_\Theta(\lambda) = mult_0(\Theta + \lambda)(mod 2) \quad (2)$$

Consider the theta function with characteristics $\lambda', \lambda'' \in \frac{1}{2}\mathbf{Z}^g$

$$\theta \begin{bmatrix} \lambda' \\ \lambda'' \end{bmatrix} (z, \tau) = \sum_{n \in \mathbf{Z}^g} \exp(\pi i(n + \lambda')\tau^t(n + \lambda') + 2\pi i(n + \lambda'')^t(z + \lambda''))$$

Then

$$\theta \begin{bmatrix} \lambda' \\ \lambda'' \end{bmatrix} (-z, \tau) = \theta \begin{bmatrix} -\lambda' \\ -\lambda'' \end{bmatrix} (z, \tau) = (-1)^{4\lambda'^t \lambda''} \theta \begin{bmatrix} \lambda' \\ \lambda'' \end{bmatrix} (z, \tau)$$

Thus if Θ is the divisor of the theta function $\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (z, \tau)$ then for any $\lambda = \lambda'\tau + \lambda'' \in A_2$ one has

$$q_\Theta(\lambda) = 4\lambda^t \lambda''(mod 2) \quad (3)$$

For any symmetric theta divisor Θ of A the bilinear form associated with q_Θ is e_2 , i.e.

$$\Delta_\lambda \Delta_\mu q_\Theta(\xi) = q_\Theta(\xi + \lambda + \mu) - q_\Theta(\xi + \lambda) - q_\Theta(\xi + \mu) + q_\Theta(\xi)$$

is independent of ξ and equals $e_2(\lambda, \mu)$. In particular one has the following formula

$$q_\Theta(\lambda + \alpha) = q_\Theta(\lambda) + e_2(\lambda, \alpha) + q_\Theta(\alpha) - q_\Theta(0) \quad (4)$$

The map $\Theta \longmapsto q_\Theta$ gives a bijective correspondence between the set of symmetric theta divisors of A and the set of quadratic forms on A_2 whose associated bilinear form equals e_2 and which vanish on $2^{g-1}(2^g + 1)$ points of A_2 .

Let $\tilde{C}$ be an algebraic, projective, smooth, irreducible curve. Let $\sigma : \tilde{C} \longrightarrow \tilde{C}$ be an involution without fixed points, let $C = \tilde{C}/\sigma$ and let $\pi : \tilde{C} \longrightarrow C$ be the factor map. Let $g = g(C), \tilde{g} = g(\tilde{C}) = 2g - 1$. We suppose that $g \geq 2$. Let $\tilde{J} = J(\tilde{C}), J = J(C)$ be the Jacobians of $\tilde{C}, C$ respectively and let $P = P(\tilde{C}, \sigma) = (1 - \sigma)\tilde{J}$ be the Prym variety. One has the maps $\pi^* : J \longrightarrow \tilde{J}, Nm : \tilde{J} \longrightarrow J$ such that $\pi^* \circ Nm = 1 + \sigma$. We denote by $j : P \longrightarrow \tilde{J}$ the embedding. The kernel of $\pi^* : J \longrightarrow \tilde{J}$ is $\{0, \eta\}$ where $\eta \in J_2$.

Conversely, given C and $\eta \in J(C)_2, \eta \neq 0$ one constructs a double unramified covering $\pi : \tilde{C} \longrightarrow C$ such that $\pi_* \mathcal{O}_{\tilde{C}} \simeq \mathcal{O}_C \oplus \mathcal{O}_C(\eta)$ and gets the set-up above.

Let $J \simeq \mathbf{C}^g/\Lambda, \tilde{J} \simeq \mathbf{C}^{2g-1}/\tilde{\Lambda}, P \simeq \mathbf{C}^{g-1}/\Lambda_-$. Choosing as in [F], [C] symplectic bases $\{a_0, \dots, a_{g-1}, b_0, \dots, b_{g-1}\}, \{\tilde{a}_0, \dots, \tilde{a}_{2g-2}, \tilde{b}_0, \dots, \tilde{b}_{2g-2}\}, \{\tilde{a}_1 - \tilde{a}_g, \dots, \tilde{a}_{g-1} - \tilde{a}_{2g-2}, \tilde{b}_1 - \tilde{b}_g, \dots, \tilde{b}_{g-1} - \tilde{b}_{2g-2}\}$ of $\Lambda, \tilde{\Lambda}$ and Λ_- respectively we can assume that $\eta = \frac{1}{2}a_0(mod \Lambda)$. Let us denote by $\tau \in \mathcal{H}_g, \tilde{\tau} \in \mathcal{H}_{2g-1}, \Pi \in \mathcal{H}_{g-1}$ the corresponding period matrices. One has the following formulas :

$$\begin{aligned} (\pi^*)_* \begin{bmatrix} \alpha_0 & \alpha \\ \beta_0 & \beta \end{bmatrix} &= \begin{bmatrix} \alpha_0 & \alpha & \alpha \\ 2\beta_0 & \beta & \beta \end{bmatrix} \\ Nm_* \begin{bmatrix} \alpha_0 & \alpha & \alpha' \\ \beta_0 & \beta & \beta' \end{bmatrix} &= \begin{bmatrix} 2\alpha_0 & \alpha + \alpha' \\ \beta_0 & \beta + \beta' \end{bmatrix} \\ j_* \begin{bmatrix} \alpha \\ \beta \end{bmatrix} &= \begin{bmatrix} 0 & \alpha & -\alpha \\ 0 & \beta & -\beta \end{bmatrix} \end{aligned} \tag{5}$$

Here $\alpha, \alpha', \beta, \beta' \in \mathbf{R}^{g-1}, \alpha_0, \beta_0 \in \mathbf{R}$ and $(\pi^*)_*, Nm_*, j_*$ are the linear maps which induce the homomorphisms π^*, Nm, j .

Wirtinger's theorem [F] states that there is a symmetric theta divisor Θ_0 of $\tilde{J}$ which is equal to $W_{\tilde{g}-1} - \pi^*\Delta$ for a certain theta characteristic Δ of C , such that $\Theta_0|_P = 2\Xi$ for some symmetric theta divisor Ξ of P . Moreover any point $L - \pi^*\Delta$ of $\Theta_0 \cap P$ satisfies the properties

$$Nm(L) \equiv K_C, h^0(\tilde{C}, L) \equiv 0(mod 2), h^0(\tilde{C}, L) \geq 2$$

Θ_0 is the divisor of the theta function $\theta[\lambda](z, \tilde{\tau})$, where

$$\lambda = \frac{1}{2}\tilde{a}_0 = \begin{bmatrix} 0 & 0 & 0 \\ \frac{1}{2} & 0 & 0 \end{bmatrix}$$

Next we recall and state in more general form some results of Welters [W]. Let $f : X \longrightarrow Y$ be a double covering of smooth, projective curves. It might be ramified. Let $\Lambda = |D|$ be a

complete linear system on Y and let $\deg(D) = d$. One denotes by S the subscheme of $X^{(d)}$ which is the pull-back of Λ under $f^{(d)}$

$$\begin{array}{ccc} S & \longrightarrow & X^{(d)} \\ \downarrow & & \downarrow \\ \Lambda & \longrightarrow & Y^{(d)} \end{array}$$

The arguments on pp.103-107 of [W] combined with Riemann-Roch's theorem give the following proposition

Proposition 0.1 *Let $\hat{D}$ be a closed point of S and let A be the maximal effective divisor of C such that $\hat{D} = \pi^*A + E$ with $A \geq 0, E \geq 0$. Let $D = f^{(d)}(\hat{D}) = 2A + E_1$. Then*

(i) *S is nonsingular at $\hat{D}$ if and only if*

$$h^0(D - A) = h^0(D) - \deg(A)$$

(ii) *Suppose S is nonsingular at $\hat{D}$. Then $f^{(d)}|_S: S \longrightarrow \Lambda$ is nondegenerate at $\hat{D}$ if and only if $f^{(d)}: X^{(d)} \longrightarrow Y^{(d)}$ is nondegenerate at $\hat{D}$ and this is the case if and only if D contains no branch points of f and $A = 0$.*

Let $B \subset Y$ be the branch locus of f and let δ be the invertible sheaf with $\delta^{\otimes 2} \simeq \mathcal{O}_Y(B)$ which determines the covering by $f_*\mathcal{O}_X \simeq \mathcal{O}_Y \oplus \delta$. An effective divisor E of X is called f -simple if $E \not\geq f^*(y)$ for any $y \in Y$. The following lemma is due to Mumford [M2].

Lemma 0.1 *Let A be a divisor of Y and let E be an effective f -simple divisor of X . Then there is an exact sequence*

$$0 \longrightarrow \mathcal{O}_Y(A) \longrightarrow f_*\mathcal{O}_X(\pi^*A + E) \longrightarrow \mathcal{O}_Y(A + Nm_f(E)) \otimes \delta^{-1} \longrightarrow 0$$

Corollary 0.1 *Under the assumptions of the preceding lemma suppose that $\deg(A) + \deg(E) < \deg(\delta)$. Then*

$$h^0(X, \pi^*A + E) = h^0(Y, A)$$

1 Double unramified coverings of elliptic curves

Let $\tilde{E}$ be an elliptic curve. Choosing a point $o \in \tilde{E}$ we shall sometimes identify $\tilde{E}$ with $J(\tilde{E})$ by the map $x \mapsto cl(x - o)$. Let $\sigma : \tilde{E} \longrightarrow \tilde{E}$ be an involution without fixed points.

Lemma 1.1 *There exists $\mu \in J(\tilde{E})$ of order 2 such that $\sigma(x) = x + \mu$. Furthermore $Ker(\pi_* : J(\tilde{E}) \longrightarrow J(E)) = \{0, \mu\}$.*

Proof. Let $\mu = \sigma(0) - 0$. Since $P(\tilde{E}, \sigma) = (\sigma - 1)J(\tilde{E}) = 0$ we have $\sigma(x - o) \equiv x - o$, thus $\sigma x = x + \mu$. Furthermore $2(\sigma(o) - o) = (1 - \sigma)(\sigma(o) - o) \equiv 0$ and $\pi_*(\sigma(o) - o) \equiv 0$, thus $Ker \pi_* = \{0, \mu\}$ since $\#Ker \pi_* = 2$. q.e.d.

Using the notation of the Introduction let $\tilde{E} \simeq \mathbf{C}/\Lambda_\tau$ where $\Lambda_\tau = \mathbf{Z}\tau + \mathbf{Z}$ with $Im(\tau) > 0$ and $\mu = \frac{1}{2}(\tau + 1)(mod \Lambda_\tau)$. Consider $\lambda(\tau)$ defined by Eq. (1). Then $\{\lambda(\tau), 1/\lambda(\tau)\}$ is the set of the roots of the equation $x^2 - k(\tau)x + 1 = 0$ where

$$k(\tau) = -(\theta_{10}(0, \tau)^8 + \theta_{01}(0, \tau)^8) / \theta_{10}(0, \tau)^4 \theta_{01}(0, \tau)^4$$

Proposition 1.1 *The map*

$$k : \Gamma_{1,2} \backslash \mathcal{H} \longrightarrow \mathbf{C} - \{0, 2\} \tag{6}$$

is an analytic isomorphism.

Proof. Let Γ_2 be the level 2 subgroup of $PSL(2, \mathbf{Z})$

$$\Gamma_2 = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} (mod 2) \right\}$$

Then $|\Gamma_{1,2} : \Gamma_2| = 2$ and the element $S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, belongs to $\Gamma_{1,2} \backslash \Gamma_2$. It is well-known (see e.g. [C]) that the map

$$\lambda : \Gamma_2 \backslash \mathcal{H} \longrightarrow \mathbf{C} - \{0, 1\}$$

given by Eq. (1) is an isomorphism. We have $S(\tau) = -1/\tau$ and $\lambda(-1/\tau) = 1/\lambda(\tau)$. The factor of $\Gamma_2 \backslash \mathcal{H}$ by the action of S is $\Gamma_{1,2} \backslash \mathcal{H}$, thus k is an analytic isomorphism. q.e.d.

Explicitly, given $k \neq 0, 2$ we find λ such that $\lambda + 1/\lambda = k$ and the corresponding pair $(\tilde{E}, \mu \in J(\tilde{E})_2)$ is given by the equation $y^2 = x(x - 1)(x - \lambda)$ and the point $\mu = cl(p_1 - p_2)$ where $p_1 = (0, 0), p_2 = (1, 0)$.

2 Extended Prym data

Let $\tilde{C}, C$ etc. be as in Section (0). Let Θ_0 be the divisor of the theta function

$$\theta \begin{bmatrix} 0 & 0 & 0 \\ \frac{1}{2} & 0 & 0 \end{bmatrix} (z, \tilde{\tau})$$

Let us denote by q_0 the quadratic form $q_{\Theta_0} : \tilde{J}_2 \longrightarrow \mathbf{Z}_2$ defined by Eq. (2). By Eq. (4) and (3) one has

$$q_0 \left(\begin{bmatrix} \alpha \\ \beta \end{bmatrix} \right) = 4\alpha^t\beta + 2\alpha_0 \pmod{2} \quad (7)$$

Hence $q_0(\rho) = 0$ for any $\rho \in P_2$. This follows also from Wirtinger's theorem. The same property holds for any symmetric theta divisor of the orbit $\{\Theta_0 + \rho \mid \rho \in P_2\}$. Let us denote $\pi^*(J)$ by B . By Eq. (5) one has $B_2 \supset P_2$.

Lemma 2.1 *A symmetric theta divisor $\Theta \subset \tilde{J}$ has the property that q_Θ vanishes on P_2 if and only if $\Theta = \Theta_0 + \alpha$ where $\alpha \in B_2$ and $q_0(\alpha) = 0$.*

Proof. Suppose $q_\Theta(P_2) = 0$. Let $\Theta = \Theta_0 + \alpha$ for some $\alpha \in \tilde{J}_2$. Then for $\rho \in P_2$ one has by Eq. (4)

$$q_\Theta(\rho) = q_0(\rho + \alpha) = q_0(\rho) + e_2(\rho, \alpha) + q_0(\alpha) - q_0(0)$$

Setting $\rho = 0$ we conclude that $q_0(\alpha) = 0$. Thus $q_\Theta(P_2) = 0$ implies that $e_2(P_2, \alpha) = 0$. Eq. (5) show that the latter holds if and only if $\alpha \in B_2$. Conversely, if $\alpha \in B_2$ and $q_0(\alpha) = 0$, then $q_\Theta(P_2) = 0$ by the formula for q_Θ above. q.e.d.

The zeros of q_0 which belong to B_2 are the following three cosets with respect to P_2 : $P_2, \lambda_1 + P_2, \lambda_2 + P_2$ where $\lambda_1 = \frac{1}{2}\tilde{a}_0 \pmod{\tilde{\Lambda}}, \lambda_2 = \frac{1}{2}\tilde{a}_0 + \frac{1}{2}\tilde{b}_0 \pmod{\tilde{\Lambda}}$. Let $\Theta_1 = \Theta_0 + \lambda_1, \Theta_2 = \Theta_0 + \lambda_2$. We conclude that there are three P_2 - orbits of symmetric theta divisors Θ such that q_Θ vanishes on P_2 :

$$\{\Theta_0 + \rho\}, \quad \{\Theta_1 + \rho\}, \quad \{\Theta_2 + \rho\}$$

These are respectively the divisors of the theta functions

$$\theta \begin{bmatrix} 0 & \alpha & \alpha \\ \frac{1}{2} & \beta & \beta \end{bmatrix} (z, \tilde{\tau}), \quad \theta \begin{bmatrix} 0 & \alpha & \alpha \\ 0 & \beta & \beta \end{bmatrix} (z, \tilde{\tau}), \quad \theta \begin{bmatrix} \frac{1}{2} & \alpha & \alpha \\ 0 & \beta & \beta \end{bmatrix} (z, \tilde{\tau})$$

where $\alpha, \beta \in \frac{1}{2}\mathbf{Z}^{g-1}$.

Lemma 2.2 *Let $\mu = \frac{1}{2}\tilde{b}_0(\text{mod } \tilde{\Lambda})$. Then $\text{Ker}(Nm) = P \cup P_\mu$, $\text{Ker}(Nm) \cap B_2 = P_2 \cup (\mu + P_2)$ and for any $x \in \tilde{C}$ there exists a unique $\xi \in P$ such that $\sigma x - x \equiv \mu + \xi$.*

Proof. It is well-known [F],[M2] that $\text{Ker}(Nm)$ has two connected components P and P_- . Since $\mu \in B_2 \setminus P_2$ and $Nm(\mu) = 0$ we get that $P_- = P_\mu$. Hence $(P \cup P_-) \cap B_2 = P_2 \cup (\mu + P_2)$. The last statement of the lemma follows from the equality $\sigma x - x + P = P_-$ [M2]. q.e.d.

Let $\tilde{J}_{2g-2}, J_{2g-2}$ be the divisor classes of degree $2g - 2$ on $\tilde{C}, C$ respectively and let $Nm : \tilde{J}_{2g-2} \longrightarrow J_{2g-2}$ be the norm map. The subvariety $Nm^{-1}(K_C + \eta)$ is a principal homogeneous space for $Nm^{-1}(0) = P \cup P_-$, thus it has two connected components. Let

$$Z = \{L \in \tilde{J}_{2g-2} \mid Nm(L) = K_C + \eta, h^0(L) \geq 1\} \quad (8)$$

and let $Z = Z_1 \cup Z_2$ where Z_i are the intersections of Z with the connected components of $Nm^{-1}(K_C + \eta)$.

Lemma 2.3 *Let M be an effective divisor of $\tilde{C}$ such that $Nm(M) \in |K_C + \eta|$. Suppose $x \in \tilde{C}$ and $x \notin Bs \mid M \mid$. Then*

$$h^0(M + \sigma x - x) = h^0(M) - 1$$

Proof. By Riemann-Roch's theorem x is a base point of $|K_{\tilde{C}} - M + x|$. Now, $K_{\tilde{C}} - M \equiv \sigma(M)$, thus $\sigma(x)$ is a base point of $|M + \sigma(x)|$ which proves the lemma since $x \notin Bs \mid M \mid$. q.e.d.

Proposition 2.1 *The theta divisors Θ_1 and Θ_2 do not contain P . The restrictions $\Theta_i.P$ are connected and reduced divisors of P which belong to the linear system $|2\Xi|$. Furthermore, up to a possible reordering of Z_i one has*

$$\Theta_i.P = Z_i - \pi^*\Delta - \lambda_i$$

The point $L - (\pi^\Delta + \lambda_i)$ is nonsingular if and only if $h^0(\tilde{C}, L) = 1$. The corresponding tangent hyperplane is equal to $Nm(|L|) \in |K_C \otimes \eta|$ via the identification*

$$T_0(P)^* \simeq H^0(\tilde{C}, K_{\tilde{C}})^- \simeq H^0(C, K_C \otimes \eta)$$

Proof. Let $\kappa_i = \pi^* \Delta + \lambda_i, i = 1, 2$. We have $Nm(\lambda_i) = \eta$, thus $Nm(\kappa_i) = K_C + \eta$ and for L with $h^0(L) \geq 1$ one has $L - \kappa_i \in \Theta_i \cap Ker(Nm)$ if and only if $Nm(|L|) \in |K_C + \eta|$. Since $dim |K_C + \eta| = g - 2$ we conclude that neither P nor P_- are contained in Θ_i . Furthermore Θ_i are ample, so $\Theta_i \cap P$ are not empty and $\Theta_i.P$ are connected divisors of P . Upon a possible reordering of Z_1 and Z_2 we have $Z_1 - \kappa_1 = \Theta_1 \cap P, Z_2 - \kappa_1 = \Theta_1 \cap P_-$. Since $P_- = P + \mu$ and $\lambda_2 = \lambda_1 + \mu$ we obtain $Z_2 - \kappa_2 = \Theta_2 \cap P$.

Claim. *For every irreducible component T of any $\Theta_i.P$ the general element $L - \kappa_i \in T$ satisfies $h^0(\tilde{C}, L) = 1$.*

Proof. Suppose the contrary. Then for any $x \in \tilde{C}$ the image of the map

$$\psi : T \times \tilde{C} \longrightarrow Ker(Nm), \quad \psi(L, x) = L + \sigma x - x$$

is contained in Z . This image must be of dimension $dim T + 1 = g - 1$. Indeed, if $M = L + \sigma x - x$, then for every sufficiently general $L \in T$ and $x \in \tilde{C}$ one has by Lemma (2.3) that $h^0(M) = h^0(L) - 1$. If $dim \psi(T \times \tilde{C}) \leq dim T$, then for every sufficiently general $M \in Im(\psi), x \in \tilde{C}$ one has $h^0(M - \sigma x + x) = h^0(L) = h^0(M) + 1$ which is an absurd by Lemma (2.3). Now, $dim Z = g - 2$, thus it is impossible that $dim \psi(T \times \tilde{C}) = g - 1$. q.e.d.

Now, suppose that $L - \kappa_i \in \Theta_i \cap P$ is an element with $h^0(L) = 1$ and let $D = |L|$. Since $Nm(D) \in |K_C + \eta|$ there is an anti invariant holomorphic differential ω of $\tilde{C}$ whose divisor of zeros is $\pi^* D$. Since $h^0(L) = 1$ the point $L - \kappa_i$ is nonsingular on Θ_i and the tangent space in $T_0 \tilde{J}$ is given by the equation $\omega = 0$. We see that Θ_i and P intersect transversely at $L - \kappa_i$ and the tangent hyperplane of $\Theta_i.P$ at $L - \kappa_i$ is given by the same equation $\omega = 0$ in $T_0 P$ since ω is anti invariant. What we have proved implies also that $Sing(\Theta_i.P) = P \cap Sing \Theta_i$. This concludes the proof of the proposition. q.e.d.

Corollary 2.1 *All irreducible components of Z are of dimension $g - 2$.*

We see that the P_2 -orbit $\{\Theta_0 + \rho\}$ is distinguished among the three P_2 -orbits of symmetric theta divisors Θ which satisfy $q_\Theta|_{P_2} = 0$ by the property that the restriction of any $\Theta_0 + \rho$ on P is equal to twice a theta divisor of P . It is also distinguished by the property that every $\{\Theta_0 + \rho\}$ contains $P_- = P + \mu$. Indeed, the fact that $\Theta_0 \supset P + \mu$ follows from the parity lemma [T] and is well-known [M1],[F]. If $\Theta_1 + \rho$ contained $P_- = P + \mu$, then Θ_2 would

contain P since $\Theta_2 = \Theta_1 + \mu$ and $\rho \in P_2$ which contradicts Proposition (2.1). Notice that the latter distinction of $\{\Theta_1 + \rho\}$ and $\{\Theta_2 + \rho\}$ parallels the distinction of $\theta_{10}(z, \tau)$ and $\theta_{01}(z, \tau)$ in the elliptic case. We can now state our extension of the Prym data.

EXTENDED PRYM DATA. *One associates to every algebraic, smooth, irreducible, projective curve $\tilde{C}$ of genus ≥ 3 with an involution $\sigma : \tilde{C} \longrightarrow \tilde{C}$ without fixed points, the principally polarized Prym variety (P, Ξ) and the two P_2 -orbits $O_1, O_2 \subset |2\Xi|$ which consist of the restrictions $\Theta.P$ of the symmetric theta divisors $\Theta \subset J(\tilde{C})$ such that $q_\Theta|_{P_2} = 0$ and $\Theta \not\subset P_-$ where $\text{Ker}(Nm_\pi : J(\tilde{C}) \longrightarrow J(C)) = P \cup P_-$*

We can now state our result which is a kind of generalization of Proposition (1.1) to curves of genus > 1 .

Theorem 2.1 *The pair $(\tilde{C}, \sigma)$ is uniquely determined up to isomorphism by the extended Prym data $(P(\tilde{C}, \sigma), \Xi), O_1, O_2 \subset |2\Xi|$.*

3 The semicanonical map and the Gauss map

In this section $K_C \in \text{Pic}^{2g-2}(C)$ is the canonical sheaf of C and $\eta \in \text{Pic}^0(C)_2$ is the sheaf with $\eta^{\otimes 2} \simeq \mathcal{O}_C$ such that $\pi_*\mathcal{O}_{\tilde{C}} \simeq \mathcal{O}_C \oplus \eta$. We shall denote by $\varphi_K, \varphi_{K \otimes \eta}$ the canonical, respectively semicanonical map of the curves under consideration. Let $L = K_C \otimes \eta$. The following lemma follows elementary from Riemann-Roch's theorem.

Lemma 3.1 *Suppose $g(C) \geq 2$. Then $|K_C \otimes \eta|$ has base points if and only if C is hyperelliptic and $\eta \simeq \mathcal{O}_C(p_1 - p_2)$ where p_1, p_2 are ramification points for the double covering $f : C \longrightarrow \mathbf{P}^1$. In this case $p_1 + p_2 = Bs |K_C \otimes \eta|$ and $K_C \otimes \eta \simeq (f^*\mathcal{O}_{\mathbf{P}^1}(g-2))(p_1 + p_2)$.*

Lemma 3.2 *Suppose $g(C) \geq 3$ and $|K_C \otimes \eta|$ is without base points. Then*

$\varphi_L : C \longrightarrow |K_C \otimes \eta|^ = \mathbf{P}^{g-2}$ is a birational embedding except in the following two cases :*

(i) $g(C) = 3$. Then $\varphi_L : C \longrightarrow \mathbf{P}^1$ is of degree 4.

(ii) $g(C) \geq 4$, C is bi-elliptic, i.e. it is a double covering $f : C \longrightarrow E$ of an elliptic curve, and $\eta \simeq f^(\epsilon)$ where $\epsilon \in \text{Pic}^0(E)_2$. Here $\varphi_L = \varphi_{\delta \otimes \epsilon} \circ f$ where δ is the invertible sheaf of E which determines the covering, i.e. $\delta^{\otimes 2} \simeq \mathcal{O}_E(x_1 + \dots + x_{2g-2})$ for the branch points $x_1, \dots, x_{2g-2}$ and $f_*\mathcal{O}_C \simeq \mathcal{O}_E \oplus \delta$.*

Proof. The case $g(C) = 3$ is clear, so let us suppose that $g \geq 4$. Let $X = \varphi_L(C)$ and let d be the degree of the map $\varphi_L : C \longrightarrow X$. We have $d \cdot \deg(X) = 2g - 2$ and $\deg(X) \geq g - 2$. This implies that the case $d \geq 3$ may occur only if $g = 4, \deg(X) = 2, d = 3$. Otherwise either φ_L is a birational embedding or $d = 2, \deg(X) = g - 1$. In the latter case $p_a(X) = 1$. Suppose $d = 2$ and X is singular. Then the normalization of X is $\hat{X} \simeq \mathbf{P}^1$ and we can decompose φ_L as

$$\varphi_L = g \circ f : C \longrightarrow \hat{X} \longrightarrow \mathbf{P}^{g-2}$$

Since φ_L is obtained from a complete linear system, g must have the same property, thus $g^* \mathcal{O}_{\mathbf{P}^{g-2}}(1) \simeq \mathcal{O}_{\mathbf{P}^1}(g - 2)$. This is impossible since $\deg(X) = g - 1$. Consequently if $d = 2$ then $X \subset \mathbf{P}^{g-2}$ is an elliptic curve. Let $E = X, f = \varphi_L : C \longrightarrow E$. Since $K_C \simeq f^*(\delta)$ and $K_C \otimes \eta \simeq f^* \mathcal{O}_E(1)$ we conclude that $\eta \simeq f^*(\epsilon)$ for some $\epsilon \in \text{Pic}^0(E)$. The covering f is ramified, so $f^* : \text{Pic}^0(E) \longrightarrow \text{Pic}^0(C)$ is an injection, hence $\epsilon \in \text{Pic}^0(E)_2$ and $\epsilon \not\simeq \mathcal{O}_E$. Conversely, if $f : C \longrightarrow E$ is a double covering of an elliptic curve, and $\eta = f^*(\epsilon)$ with $\epsilon \in \text{Pic}^0(E)_2, \epsilon \not\simeq \mathcal{O}_E$ then

$$H^0(C, K_C \otimes \eta) \simeq H^0(C, \pi^*(\delta \otimes \epsilon)) = \pi^* H^0(E, \delta \otimes \epsilon)$$

Hence $\varphi_L = \varphi_{\delta \otimes \epsilon} \circ f$ and $d = 2$.

It remains to rule out the possibility $g = 4, d = 3, \deg(X) = 2$. Here $f = \varphi_L : C \longrightarrow X \simeq \mathbf{P}^1$ so $L \simeq M^{\otimes 2}$, where $\deg(M) = 3, h^0(C, M) = 2$, and $|M|$ is without base points. Thus C is not hyperelliptic and

$$\varphi_K(C) = Q \cap F \subset \mathbf{P}^3$$

where Q is a quadric and F is a cubic surface. We have $|L| = |M| + |M|$ since $\dim |L| = 2$. Let l_1, l_2 be lines in Q such that $l_1 + l_2 = Q.H$ for a plane H . Let $M_i = \mathcal{O}_C(C.l_i)$. We can assume that $M = M_1$. Then $\eta \simeq M_1 \otimes M_2^{-1}$. If Q were singular, then $M_1 \simeq M_2$, so $\eta \simeq \mathcal{O}_C$ which is absurd. Suppose Q is nonsingular. Then $\eta^{\otimes 2} \simeq \mathcal{O}_C$ implies $|M_2^{\otimes 2}| = |M_1^{\otimes 2}| = |M_1| + |M_1|$. This is again impossible since any reduced divisor $x_1 + x_2 + x_3 \in |M_2|$ can have only two common points with any two divisors $D_1, D_2 \in |M_1|$. q.e.d.

Suppose $g \geq 3$. Following Welters [W] let S be the subscheme of $\tilde{C}^{2g-2}$ which is the pull-back of $|K_C \otimes \eta| \subset C^{2g-2}$

$$\begin{array}{ccc} S & \longrightarrow & \tilde{C}^{2g-2} \\ \downarrow & & \downarrow Nm \\ |K_C \otimes \eta| & \longrightarrow & C^{2g-2} \end{array}$$

It breaks naturally into two disjoint subschemes $S = S_1 \cup S_2$. The singularities of S can be calculated by Proposition (0.1) with $X = \tilde{C}, Y = C, f = \pi, \pi^{(2g-2)} = Nm$. Since S is a locally complete intersection and $\pi^{(2g-2)}$ is a finite map every irreducible component of S has dimension $g - 2$. A Zariski open, dense subset of $|K_C \otimes \eta|$ consists of reduced divisors by Lemmas (3.1) and (3.2), so according to Proposition (0.1) S is reduced. The subvarieties S_1, S_2 are connected provided $g \geq 3$ [W].

Let T_1, T_2 be divisors from $|2\Xi|$ which belong to the orbits O_1, O_2 respectively. Suppose $g \geq 3$. The Gauss maps $G_i : T_i^{ns} \longrightarrow \mathbf{P}(T_0P)^*$ are defined on the nonsingular loci of T_i and send a point $x \in T_i^{ns}$ to the translation of the tangent hyperplane $T_x(T_i)$ to $0 \in P$. Let $T = T_1 \sqcup T_2$ and let $G : T^{ns} \longrightarrow \mathbf{P}(T_0P)^*$ be the map whose restriction on T_i^{ns} equals G_i . Let S^0 be the Zariski open subset of S which consists of those $\hat{D}$ with $h^0(\tilde{C}, \hat{D}) = 1$. By Proposition (2.1) the map $cl : S^0 \longrightarrow Z^{ns}$ is an isomorphism and moreover one can identify T_i^{ns} with Z_i^{ns} by translation and $\mathbf{P}(T_0P)^*$ with $|K_C \otimes \eta|$. Since the Gauss map does not depend on the translation one has for the Gauss map $G : Z^{ns} \longrightarrow |K_C \otimes \eta|$ and every $\hat{D} \in S^0$ the formula

$$G(cl(\hat{D})) = Nm(\hat{D}) \tag{9}$$

Proposition 3.1 *Let $L = K_C \otimes \eta$. Suppose $g \geq 4$. Let $R \subset T_i$ be the ramification locus of G and let B be the algebraic closure of $G(R)$.*

(i) *If C is hyperelliptic and $\eta \simeq \mathcal{O}_C(p_1 - p_2)$, where p_1, p_2 are Weierstrass points of C , then $P(\tilde{C}, \sigma) \simeq J(C_2)$ for a certain hyperelliptic curve C_2 (see Section (4)) and*

$$B = \varphi_K(C_2)^* \cup \bigcup_{i=1}^{2g} \varphi_K(q_i)^*$$

where $\varphi_K(C_2)^$ is the dual hypersurface of $\varphi_K(C_2)$ and $\varphi_K(q_i)^*$ are the stars of hyperplanes which contain $\varphi_K(q_i)$, where q_i are the Weierstrass points of C_2 .*

(ii) If $|K_C \otimes \eta|$ is without base points and $\varphi_L : C \longrightarrow |K_C \otimes \eta|^*$ is a birational embedding, then B has a unique irreducible component of dimension $g - 3$ and degree > 1 . This component is equal to $\varphi_L(C)^*$.

(iii) If $|K_C \otimes \eta|$ is without base points and

$$f = \varphi_L : C \longrightarrow \varphi_L(C) = E$$

is a map of degree 2 onto an elliptic curve, then

$$B = E^* \cup \bigcup_{i=1}^{2g-2} x_i^*$$

where x_i are the branch points of φ_L .

Proof. (i) In this case any T_i is the union of two translates of the theta divisor $\Xi \subset P$ as it will be shown in Section (4). So, Part(i) follows from the description of the branch locus of the Gauss map of the theta divisor of a hyperelliptic Jacobian [A].

Now, let us assume that $|K_C \otimes \eta|$ is without base points. If $Nm : S^0 \longrightarrow |K_C \otimes \eta|$ is degenerate at $\hat{D} \in S^0$, then by Proposition (0.1) $\hat{D} = \pi^*A + E$ for some $A > 0$, so $D = Nm(\hat{D}) = 2A + Nm(E)$. Let $H \subset |K_C \otimes \eta|^*$ be the hyperplane which corresponds to D . Either H contains an image of a ramification point of the map $\varphi_L : C \longrightarrow |K_C \otimes \eta|^*$, or H is tangent to a branch $\varphi_L(U)$ at a point $\varphi_L(p)$, where $p \in U \subset C$ and φ_L is nondegenerate at p . The former case can happen only for finitely many points. This shows that any component of B of dimension $g - 3$ must be either a star of hyperplanes which contain a branch point $\varphi_L(p)$, or it is contained in the dual variety $\varphi_L(C)^*$.

Proof of (ii). It remains to show that $\varphi_L(C)^* \subset B$. Let the hyperplane $H \subset |K_C \otimes \eta|^*$ be a sufficiently general element of $\varphi_L(C)^*$ and let

$$D = \varphi_L^*(H) = 2p + p_3 + \dots + p_{2g-2}$$

be the corresponding divisor of $|K_C \otimes \eta|$. Here $p, p_3, \dots, p_{2g-2}$ are distinct points of C and φ_L is not degenerate at p .

Claim 1. Let $\pi^{-1}(p) = p' + p''$. One can choose $p'_i \in \tilde{C}$ with $\pi(p'_i) = p_i$ so that

$$\hat{D} = p' + p'' + p'_3 + \dots + p''_{2g-2}$$

has the property $h^0(\hat{D}) = 1$.

Proof. Let us choose arbitrary points $q_i \in \tilde{C}$ such that $\pi(q_i) = p_i$. Let

$$\hat{D}_0 = p' + p'' + q_3 + \dots + q_{2g-2}$$

If $h^0(\hat{D}) \geq 2$, then at least one of the points q_i is not a base point of $|\hat{D}_0|$. Indeed, otherwise

$$h^0(\mathcal{O}_{\tilde{C}}(\pi^*p)) = h^0(\mathcal{O}_C(p)) + h^0(\mathcal{O}_C(p) \otimes \eta) \geq 2$$

which is possible only in the Case (i). If $q_i \notin Bs|\hat{D}_0|$, let $\hat{D}_1 = \hat{D}_0 + \sigma(q_i) - q_i$. Then by Lemma (2.3) $h^0(\hat{D}_1) = h^0(\hat{D}_0) - 1$. Repeating the same argument with $\hat{D}_1$ etc. we obtain eventually the required divisor $\hat{D}$. q.e.d.

Now, $\hat{D} \in S^0$, the Gauss map at the point $cl(\hat{D}) \in Z^{ns}$ equals H by Eq. (9) and it is ramified at $\hat{D}$ according to Proposition (0.1). So, $\varphi_L(C)^* \subset B$.

Proof of (iii). We have proved above that

$$B \subset E^* \cup \bigcup_{i=1}^{2g-2} x_i^*$$

Let H be a sufficiently general element of E^* and let

$$D = \varphi_L^*(H) = 2p + 2q + p_5 + \dots + p_{2g-2}$$

be the corresponding divisor of $|K_C \otimes \eta|$. Here $p, q, p_5, \dots, p_{2g-2}$ are distinct points of C , φ_L is nondegenerate at p, q and $\{p, q\} = f^*(x)$ for some $x \in E$.

Claim 2. Let $\pi^{-1}(p) = \{p', p''\}, \pi^{-1}(q) = \{q', q''\}$. One can choose $p'_i \in \tilde{C}$ with $\pi(p'_i) = p_i$ so that

$$\hat{D} = p' + p'' + 2q' + p'_5 + \dots + p'_{2g-2}$$

has the property $h^0(\hat{D}) = 1$.

Proof. Let $\eta = f^*(\epsilon)$ and let $y \in E$ be the point such that $\epsilon \simeq \mathcal{O}_E(y - x)$. Then one has canonical isomorphisms

$$\begin{aligned} H^0(\mathcal{O}_{\tilde{C}}(\pi^*(p + q))) &\simeq \pi^*(H^0(\mathcal{O}_C(p + q)) \oplus H^0(\mathcal{O}_C(p + q) \otimes \eta)) \\ &\simeq (\pi^* \circ f^*)(H^0(\mathcal{O}_E(x)) \oplus H^0(\mathcal{O}_E(y))) \simeq \mathbf{C}^2 \end{aligned}$$

Thus $|p' + p'' + q' + q''|$ is a pencil without base points. Using the same argument as in Claim 1 we conclude that one can choose p'_i so that

$$\hat{D}' = p' + p'' + q' + q'' + p'_5 + \dots + p'_{2g-2}$$

has the property $h^0(\hat{D}') = 2$. Applying once more Lemma (2.3) we conclude that $\hat{D} = \hat{D}' + q' - q''$ satisfies $h^0(\hat{D}) = 1$. q.e.d.

We conclude as in Part (ii) that $E^* \subset B$. If $x_i = \varphi_L(q_i)$ is a branch point of f and H is a sufficiently general hyperplane in $|K_C \otimes \eta|^*$ which contains x_i , then the corresponding divisor of $|K_C \otimes \eta|$ has the form

$$D = \varphi_L^*(H) = 2q_i + p_3 + \dots + p_{2g-2}$$

where $q_i, p_3, \dots, p_{2g-2}$ are distinct. The same argument as in Part (ii) proves that $H \in B$, so $x_i^* \subset B$. Proposition (3.1) is proved. q.e.d.

We see that if $g \geq 4$ then the branch locus B of the Gauss map $G : T^{ns} \longrightarrow \mathbf{P}(T_0P)^*$ has a unique irreducible component B_0 of dimension $g - 3$ and degree ≥ 2 . We have shown above that $B = X^*$ for a certain irreducible curve $X \subset \mathbf{P}(T_0P)^*$. So, for $g \geq 4$, by the equality $(X^*)^* = X$ [H-K] we obtain that B_0^* is a curve X . The following alternative takes place.

(i) $\deg(X) = g - 2$

(ii) $\deg(X) = 2g - 2$

(iii) $\deg(X) = g - 1$

The three cases correspond to those in Proposition (3.1). If Case (ii) occurs we prove Theorem (2.1) as follows: C is isomorphic to the normalization of X . The normalization map $f : C \longrightarrow X \subset \mathbf{P}(T_0P)^*$ is associated to the complete linear system $|K_C \otimes \eta|$. Thus we obtain $\eta \in J(C)_2$.

Cases (i) and (iii) are considered respectively in Sections (4) and (5). The case $g = 3$ is treated in Section (6) and the case $g = 2$ in Section (4).

4 The hyperelliptic case, $g \geq 2$

Throughout this section we suppose that C is a hyperelliptic curve of genus $g \geq 2$, $f : C \longrightarrow \mathbf{P}^1$ is the double covering and $\eta \simeq \mathcal{O}_C(p_1 - p_2)$ where p_1, p_2 are ramification points of f and $p_1 \neq p_2$. Let $R = \{p_1, p_2, p_3, \dots, p_{2g+2}\}$ be the set of ramification points of




Figure 1:

$f, R_1 = \{p_1, p_2\}, R_2 = R \setminus R_1$. Let $B_i = f(R_i), i = 1, 2$. According to [M2],[Da] the covering $f \circ \pi : \tilde{C} \longrightarrow C \longrightarrow \mathbf{P}^1$ has Galois group $\mathbf{Z}_2 \times \mathbf{Z}_2$. In the corresponding diagram of Fig.1 $f_i : C_i \longrightarrow \mathbf{P}^1$ is branched at $B_i, i = 1, 2$. Furthermore $\pi_2^* : J(C_2) \longrightarrow P(\tilde{C}, \sigma)$ is an isomorphism. Let $\Theta_0 = W_{\tilde{g}-1}(\tilde{C}) - \pi^* \Delta$ be as in Section (2).

Lemma 4.1 *Let $\Xi \subset P(\tilde{C}, \sigma) = P$ be a symmetric theta divisor. Then there exists a unique $\rho \in P_2$ such that*

$$\Xi = \pi_1^*(\zeta_1) + \pi_2^*W_{g-2}(C_2) - \pi^*\Delta - \rho \quad (10)$$

where ζ_1 is the rational equivalence class of the points of the rational curve C_1 .

Proof. By Wirtinger's theorem there is a unique translation $\Theta = \Theta_0 + \rho$ with $\rho \in P_2$ such that $\Theta.P = 2\Xi$. The points of $\Theta \cap P$ have the form $L - \pi^*\Delta - \rho$ where $Nm(L) = K_C$, $h^0(\tilde{C}, L) \equiv 0(mod 2)$ and $h^0(\tilde{C}, L) \geq 2$. Now, $|K_C| \simeq f^*|\mathcal{O}_{\mathbf{P}^1}(g-1)|$. One easily checks that if $\hat{D}$ is effective divisor of $\tilde{C}$ and $Nm(\hat{D}) \in f^*|\mathcal{O}_{\mathbf{P}^1}(g-1)|$ then

$$\hat{D} = \pi_1^*E + \pi_2^*F$$

where E and F are effective divisors of C_1, C_2 respectively. We have $deg(E) + deg(F) = g-1$ which gives only two irreducible components of dimension $\geq g-2$ of

$$Nm^{-1}(K_C) \cap W_{2g-2}(\tilde{C})$$

namely $\pi_2^*W_{g-1}(C_2)$ of dimension $g-1$ and $\pi_1^*(\zeta_1) + \pi_2^*W_{g-2}(C_2)$ of dimension $g-2$. On the other hand the above intersection has, by the general theory, two irreducible components: a translation of Ξ and a translation of P_- . This shows Eq. (10). q.e.d.

Now, let us calculate the orbits $O_1, O_2 \subset |2\Xi|$. Let $T_i \in O_i, i = 1, 2$. By Proposition (2.1) one has

$$T_i = Z_i - \pi^*\Delta - \nu_i \quad (11)$$

for some $\nu_i \in \lambda_i + P_2, i = 1, 2$.

Lemma 4.2 *One can enumerate $\pi^{-1}(p_1)$ as $\{p'_1, p''_1\}$ and $\pi^{-1}(p_2)$ as $\{p'_2, p''_2\}$ so that*

$$\begin{aligned} Z_1 &= \pi_2^*W_{g-2}(C_2) + p'_1 + p'_2 \cup \pi_2^*W_{g-2}(C_2) + p''_1 + p''_2 \\ Z_2 &= \pi_2^*W_{g-2}(C_2) + p'_1 + p''_2 \cup \pi_2^*W_{g-2}(C_2) + p''_1 + p'_2 \end{aligned}$$

Proof. From Lemma (3.1) we have

$$|K_C \otimes \eta| = f^* | \mathcal{O}_{\mathbf{P}^1}(g-2) | + p_1 + p_2$$

One easily checks that if $\hat{D}$ is effective divisor of $\tilde{C}$ and $Nm(\hat{D}) \in f^* | \mathcal{O}_{\mathbf{P}^1}(g-2) |$ then

$$\hat{D} = \pi_1^*E + \pi_2^*F$$

where E, F are effective divisors of C_1, C_2 respectively. We have $\deg(E) + \deg(F) = g-2$. Thus the only irreducible component of dimension $\geq g-2$ of

$$Nm^{-1}(f^*\mathcal{O}_{\mathbf{P}^1}(g-2)) \cap W_{2g-4}(\tilde{C})$$

is $\pi_2^*W_{g-2}(C_2)$. The irreducible components of $Z = Z_1 \cup Z_2$ are of dimension $g-2$ by Corollary (2.1) and the transformation $L \mapsto L + \sigma(p) - p$ interchanges the two components of $Nm^{-1}(K_C \otimes \eta)$. This shows that Z_1 and Z_2 have the form given in the lemma. q.e.d.

Lemmas (4.1), (4.2) and Eq. (11) give the following corollary

Corollary 4.1 *Let Ξ be an arbitrary symmetric theta divisor of $P(\tilde{C}, \sigma)$ and let T_1, T_2 be two divisors of the orbits $O_1, O_2 \subset |2\Xi|$ respectively. Then*

$$\begin{aligned} T_1 &= \Xi + p'_1 + p'_2 - \pi_1^*(\zeta_1) - \mu_1 \cup \Xi + p''_1 + p''_2 - \pi_1^*(\zeta_1) - \mu_1 \\ T_2 &= \Xi + p'_1 + p''_2 - \pi_1^*(\zeta_1) - \mu_2 \cup \Xi + p''_1 + p'_2 - \pi_1^*(\zeta_1) - \mu_2 \end{aligned}$$

for some $\mu_i \in \lambda_i + P_2, i = 1, 2$.

Let us choose in an arbitrary way Ξ, T_1, T_2 as above and let us denote by x_1, y_1, x_2, y_2 the elements of $P(\tilde{C}, \sigma)$ such that $T_1 = \Xi + x_1 \cup \Xi + y_1, T_2 = \Xi + x_2 \cup \Xi + y_2$. There are two possible ways of representing the set $\{x_1, y_1, x_2, y_2\}$ as union $A \cup B$, where $\#A = \#B = 2$, and A, B have one point of $\{x_1, y_1\}$ and one point of $\{x_2, y_2\}$. Namely as :

$$\begin{aligned} \{x_1, x_2\} &\cup \{y_1, y_2\}, \\ \{x_1, y_2\} &\cup \{y_1, x_2\} \end{aligned} \quad (12)$$

Taking the sums of the sets in (12), using Corollary (4.1) and taking into account that $\lambda_1 + \lambda_2 = \mu$ (Section (2)) we see that the extended Prym data determines the following P_2 -orbit of quadruples of points in $P(\tilde{C}, \sigma)$, each quadruple being split into a union of two pairs:

$$\begin{aligned} &\{2p'_1 + \pi^*p_2 - \pi_1^*(2\zeta_1) - \mu + \rho, 2p''_1 + \pi^*p_2 - \pi_1^*(2\zeta_1) - \mu + \rho\} \\ &\cup \{2p'_2 + \pi^*p_1 - \pi_1^*(2\zeta_1) - \mu + \rho, 2p''_2 + \pi^*p_1 - \pi_1^*(2\zeta_1) - \mu + \rho\} \end{aligned} \quad (13)$$

where $\rho \in P_2$. The splitting of the quadruples is consistent with the action of P_2 on Q .

Let $f_2^{-1}(f(p_i)) = \{q'_i, q''_i\}$ where $\pi_2^{-1}(q'_i) = p'_i, \pi_2^{-1}(q''_i) = p''_i, i = 1, 2$. Let $f_2^{-1}(f(p_j)) = q_j$ for $3 \leq j \leq 2g + 2$. Since $\pi_2 : C \longrightarrow C_2$ is branched at $\{q'_1, q''_1, q'_2, q''_2\}$ one has

$$2p'_i = \pi_2^*(q'_i), \quad 2p''_i = \pi_2^*(q''_i) \quad (14)$$

One has also that

$$\pi_1^*(2\zeta_1) = \pi_1^*f_1^*(f(p_i)) = 2\pi^*(p_i) \quad (15)$$

Claim For any i with $1 \leq i \leq 2$ and any j with $3 \leq j \leq 2g + 2$ there is a point $\rho_{ij} \in P_2$ such that $\mu = -\pi^*(p_i - p_j) - \rho_{ij}$.

Proof. By Eq. (5) one has $\pi^*J(C)_2 = P_2 \cup (\mu + P_2)$. Since $P(\tilde{C}, \sigma) = \pi^*J(C_2)$ one concludes by the description of the points of order 2 of the hyperelliptic Jacobian $J(C_2)$ [M3] that

$$P_2 = \{\pi_2^*(S_1 - S_2) \mid S_1 \cup S_2 \subset R_2, S_1 \cap S_2 = \emptyset, \#S_1 = \#S_2\}$$

Using this one easily shows that $\pi^*(p_i - p_j) \notin P_2$. q.e.d.

Now, using Eq. (14),(15) and the Claim we have for any j with $3 \leq j \leq 2g + 2$

$$\begin{aligned} 2p'_1 + \pi^*p_2 - \pi_1^*(2\zeta_1) - \mu + \rho &= \pi_2^*(q'_1) - \pi^*(p_2) - \mu + \rho \\ &= \pi_2^*(q'_1) - \pi^*(p_j) + \rho_{2j} + \rho = \pi_2^*(q'_1 - q_j) + \rho_{2j} + \rho \end{aligned}$$

We obtain that the P_2 -orbit Q is equal to

$$\begin{aligned} &\{\pi_2^*(q'_1 - q_j) + \rho, \pi_2^*(q''_1 - q_j) + \rho\} \\ &\cup \{\pi_2^*(q'_2 - q_j) + \rho, \pi_2^*(q''_2 - q_j) + \rho\} \end{aligned} \quad (16)$$

Reconstruction of (C, η) in the hyperelliptic case, $g \geq 3$.

We have a polarized isomorphism $P(\tilde{C}, \sigma) \simeq J(C_2)$. So, by Torelli's theorem [ACGH] one reconstructs the smooth, hyperelliptic curve C_2 of genus $g_2 = g - 1 \geq 2$. It has a unique complete linear system g_2^1 . Take a point $q \in C_2$ such that $2q \in g_2^1$. Consider the Abel map $\alpha : C_2 \longrightarrow J(C_2)$ given by $\alpha(x) = cl(x - q)$.

Lemma 4.3 *There is a unique quadruple of Q whose points belong to $\alpha(C_2)$. Any other quadruple has no points in common with $\alpha(C_2)$.*

Proof. If $q = q_j$ we set $\rho = 0$ in (16) and see that the quadruple

$$\{\{\alpha(q'_1), \alpha(q''_1)\} \cup \{\alpha(q'_2), \alpha(q''_2)\}\} \quad (17)$$

is contained in $\alpha(C_2)$. Suppose that for some $\rho \in J(C_2)_2, \rho \neq 0$ one has

$$q'_1 - q_j + \rho \equiv x - q_j.$$

Multiplying by 2 both sides of this equality we obtain $2q'_1 \equiv 2x$. Since C_2 has a unique g_2^1 and $q'_1 \neq x$ for $\rho \neq 0$ we conclude that $2q'_1 \in g_2^1$ which is an absurd. This argument shows that none of the quadruples of Q different from (17) can have points in common with $\alpha(C_2)$. q.e.d.

Now, we choose a map $f_2 : C_2 \longrightarrow \mathbf{P}^1$ of degree 2 and observe that the quadruple of points of C_2 defined in the lemma is transformed by f_2 into a set of two points. Furthermore this set does not depend on the choice of the ramification point q of f_2 . Let us denote it by B_1 . Let B_2 be the branch locus of f_2 . Then C is isomorphic to the hyperelliptic curve branched at $B = B_1 \cup B_2$ and $\eta \in J(C)_2$ corresponds to this partition of B [M3].

Reconstruction of (C, η) in the case $g = 2$.

Here $P(\tilde{C}, \sigma)$ is an elliptic curve E . Let $o \in E$ be the zero, let φ_1, φ_2 be a basis of $H^0(E, \mathcal{O}_E(2o))$ and let $f_2 = (\varphi_1 : \varphi_2) : E \longrightarrow \mathbf{P}^1$. For any $\rho \in E_2$, if $t_\rho : E \longrightarrow E$ is the translation by ρ , there exists $\psi \in PGL(2)$ such that the following diagram is commutative

$$\begin{array}{ccccc} E & \xrightarrow{t_\rho} & E & & \\ f_2 \downarrow & & \downarrow f_2 & & \\ \mathbf{P}^1 & \xrightarrow{\psi} & \mathbf{P}^1 & & \end{array} \quad (18)$$




Figure 2:

Moreover ψ permutes the branch points of f_2 . Let B_2 be the branch locus of f_2 . Take any of the quadruples of Q . Each of its two pairs is invariant under the action of $-id_E$. Thus the image of the quadruple is a set of two points which we denote by B_1 . If we choose another quadruple of Q with image B'_1 , then (18) shows that there is a $\psi \in PGL(2)$ such that $\psi(B_1) = B'_1, \psi(B_2) = B_2$. This gives the reconstruction of (C, η) , up to isomorphism, as the hyperelliptic curve branched at $B = B_1 \cup B_2$ and η as the point of $J(C)_2$ which corresponds to this partition of B .

5 The bi-elliptic case, $g \geq 4$

Let $f : C \longrightarrow E$ be a double covering of an elliptic curve E ramified at $B = \{x_1, \dots, x_{2g-2}\}$ and determined by $\delta \in Pic^{g-1}(E)$ with $\delta^{\otimes 2} \simeq \mathcal{O}_E(B)$. Suppose $\eta = f^*(\epsilon)$ for some $\epsilon \in Pic^0(E)_2$. Then the unramified covering $\pi : \tilde{C} \longrightarrow C$ determined by η fits into the commutative diagram of Fig.2 where $deg(f_i) = deg(\pi_i) = 2$, $f_1 : C_1 \longrightarrow E$ is unramified, determined by ϵ , $f_2 : C_2 \longrightarrow E$ is ramified at B and is determined by $\delta_2 = \delta \otimes \epsilon$. Here we have the assumptions of Part (iii) of Proposition (3.1) so the extended Prym data determines :

- E as the curve isomorphic to the dual $X \subset \mathbf{P}(T_0P)$ of the unique irreducible component of degree ≥ 2 of the branch locus $G(R)$ of the Gauss map $G : T^{ns} \longrightarrow \mathbf{P}(T_0P)^*$.
- The points $\{x_i \mid i = 1, \dots, 2g-2\}$ as the duals of the remaining irreducible components

of $G(R)$.

- $\delta_2 \simeq \delta \otimes \epsilon$ as isomorphic to $\mathcal{O}_X(1)$.

So, it remains to reconstruct ϵ which is the content of the rest of this section.

Lemma 5.1 *Let $T_1 = Z_1 - \pi^*\Delta - \mu_1, T_2 = Z_2 - \pi^*\Delta - \mu_2$ be arbitrary divisors of the orbits $O_1, O_2 \subset |2\Xi|$, where $\mu_i \in \lambda_i + P_2$ (Section (2)). Then T_1, T_2 are irreducible. Reordering, if necessary, $\{\lambda_1, \lambda_2\}$, respectively $\{O_1, O_2\}, \{Z_1, Z_2\}, \{T_1, T_2\}$ one has that the elements $e_1 \in T_1, e_2 \in T_2$ have the form*

$$(i) \ e_1 = \pi_1^*(\xi_1) + \pi_2^*(\xi_2) - \pi^*\Delta - \mu_1$$

where $\xi_1 \in C_1, \xi_2 \in W_{g-2}(C_2)$ and $Nm_{f_1}(\xi_1) + Nm_{f_2}(\xi_2) = \delta_2$.

$$(ii) \ e_2 = \pi_2^*(\xi_2) - \pi^*\Delta - \mu_2$$

where $\xi_2 \in W_{g-1}(C_2)$ and $Nm_{f_2}(\xi_2) = \delta_2$.

Proof. One has to calculate the irreducible components of Z defined in (8). One has

$$H^0(C, K_C \otimes \eta) \simeq H^0(C, f^*\delta_2) \simeq H^0(E, \delta_2) \oplus H^0(E, \delta_2 \otimes \delta^{-1}) \simeq H^0(E, \delta_2)$$

Thus $|K_C \otimes \eta| = f^*|\delta_2|$. If $\hat{D}$ is an effective divisor of $\tilde{C}$ such that $Nm_\pi(\hat{D}) \in f^*|\delta_2|$, then

$$\hat{D} = \pi_1^*E + \pi_2^*F$$

where E and F are effective divisors of C_1, C_2 respectively. One has $Nm_\pi \circ \pi_i^* = f^* \circ Nm_{f_i}$, so

$$Nm_{f_1}(E) + Nm_{f_2}(F) \equiv \delta_2$$

Corollary (2.1) and a dimension count show that $cl(\hat{D})$ might be a general element of Z if either $deg(E) = 1, deg(F) = g - 2$ or $E = 0, deg(F) = g - 1$. So, $Z = Z' \cup Z''$ where

$$Z' = \{\pi_1^*(\xi_1) + \pi_2^*(\xi_2) \mid \xi_1 \in C_1, \xi_2 \in W_{g-2}(C_2), Nm_{f_1}(\xi_1) + Nm_{f_2}(\xi_2) \equiv \delta_2\}$$

and

$$Z'' = \{\pi_2^*(\xi_2) \mid \xi_2 \in W_{g-1}(C_2), Nm_{f_2}(\xi_2) \equiv \delta_2\}$$

Claim 1. Z' is irreducible.

Proof. We consider the map $h : C_2^{(g-2)} \longrightarrow E$ defined by $h(D) = |\delta_2 - Nm_{f_2}(D)|$ and the pull-back diagram

$$\begin{array}{ccc} X & \longrightarrow & C_1 \\ \downarrow & & \downarrow f_1 \\ C_2^{(g-2)} & \xrightarrow{h} & E \end{array}$$

Then Z' is the image of X under the map

$$(D, x) \longmapsto cl(\pi_1^*(x) + \pi_2^*(D))$$

Now, X might be reducible if $h_*\pi_1(C_2^{(g-2)})$ is contained in $f_{1*}\pi_1(C_1)$. This is impossible. Indeed, $f_{2*} : H_1(C_2) \longrightarrow H_1(E)$ is epimorphic since f_2 is ramified. Therefore

$$(cl \circ f_2^{(g-2)})_* : H_1(C_2^{(g-2)}) \longrightarrow H_1(J_{g-2}(E))$$

is epimorphic. Composing it with the isomorphism $J_{g-2}(E) \longrightarrow E$ given by $\xi \mapsto |\delta_2 - \xi|$ one obtains that $h_* : H_1(C_2^{(g-2)}) \longrightarrow H_1(E)$ is epimorphic. This proves that X and therefore Z' are irreducible. q.e.d.

Claim 2. Z'' is irreducible.

Proof. With the same notation as above one considers the pull-back diagram

$$\begin{array}{ccc} Y & \longrightarrow & C_2 \\ \downarrow & & \downarrow f_2 \\ C_2^{(g-2)} & \xrightarrow{h} & E \end{array}$$

Then Z'' is the image of Y under the map

$$(D, y) \longmapsto cl(\pi_2^*(D + y))$$

In order to prove that Y is irreducible it suffices to verify that not every component of the branch divisor $h^*(B)$ has even multiplicity. Now, h can be decomposed as

$$h = p \circ f_2^{(g-2)} : C_2^{(g-2)} \longrightarrow E^{(g-2)} \longrightarrow E$$

where p is the fiber bundle map defined by $p(A) = |\delta_2 - A|$. Let $x \in B$. Then $|\delta_2 - x|$ is a linear system of degree $g - 2 \geq 2$ without base points. Let $A = p_1 + \dots + p_{g-2}$ be an element

with no points in common with B . Let $D \in C_2^{(g-2)}$ with $f_2^{(g-2)}(D) = A$. Then $p^{-1}(x)$ is smooth at A and $f_2^{(g-2)}$ is nondegenerate at D , thus $h^*(x)$ is a divisor with multiplicity 1. q.e.d.

Now, Z has two connected components Z_1 and Z_2 , enumerated as in Proposition (2.1). So, $Z' \neq Z''$, Z_i are irreducible and either $Z_1 = Z'$, $Z_2 = Z''$ or $Z_1 = Z''$, $Z_2 = Z'$. Reordering $\{Z_1, Z_2\}$ if necessary we can assume that the former case takes place. q.e.d.

Lemma 5.2 *The singular locus of Z_1 has codimension ≥ 2 .*

Proof. Consider a divisor of $\tilde{C}$ of the form $\pi_1^*A + H$ where H is effective, π_1 -simple and $\deg(A) \geq 1$. Then by Corollary (0.1) one concludes that

$$h^0(\tilde{C}, \pi_1^*A + H) = h^0(C_1, A) \quad (19)$$

Let $\hat{D} = \pi_1^*(x) + \pi_2^*(F)$ where $x \in C_1$, F is effective divisor of C_2 and $f_1(x) + Nm_{f_2}(F) \equiv \delta_2$. Proposition (2.1) and (19) show that $\mathcal{O}_{\tilde{C}}(\hat{D})$ is a singular point of Z_1 if and only if F is not π_1 -simple. Now, if $\pi_1^*(y) \leq \pi_2^*(F)$, then one easily checks that $f_2^*(f_1(y)) \leq F$. Thus $Sing(Z_1)$ consists of

$$cl(\pi_1^*(x + f_1^*(t)) + \pi_2^*(G)) \quad (20)$$

where $x \in C_1$, $t \in E$, $G \in C_2^{(g-4)}$ and $Nm_{f_2}(G) \in |\delta_2 - f_1(x) - 2t|$. For any $G \in C_2^{(g-4)}$ there are two different $\zeta_1 \in J_3(C_1)$ such that $Nm_{f_1}(\zeta_1) \equiv \delta_2 - Nm_{f_2}(G)$. Thus the elements of the type (20) form a sublocus of Z_1 of dimension $\leq g - 4$. q.e.d.

Lemma 5.3 *$Sing(Z_2)$ has a unique irreducible component V of codimension 1 in Z_2 . A Zariski open, dense subset of V consists of the elements*

$$cl(\pi_2^*(f_2^*(x) + G)) \quad (21)$$

where $x \in E$, and G is an effective, f_2 -simple divisor of C_2 such that $Nm_{f_2}(G) \in |\delta_2 - 2x|$.

Proof. We claim that

$$\dim W_{g-1}^1(C_2) \cap Nm_{f_2}^{-1}(\delta_2) \leq g - 4 \quad (22)$$

This is clear if C_2 were not hyperelliptic. If C_2 were hyperelliptic, then $W_{g-2}^1(C_2) = g_2^1 + W_{g-3}(C_2)$. This irreducible variety can not be contained in $Nm_{f_2}^{-1}(\delta_2)$. Indeed, otherwise

its translation would be contained in the abelian hypersurface $Nm_{f_2}^{-1}(0)$ of $J(C_2)$ which is absurd since this translation generates $J(C_2)$. By (22) we conclude that the sublocus of $Sing(Z_2)$:

$$\{\pi_2^*(\xi_2) \mid Nm_{f_2}(\xi_2) = \delta_2, h^0(C, \xi_2) \geq 2\} \quad (23)$$

has codimension ≥ 2 in Z_2 .

Suppose F is an effective, f_2 -simple divisor of C_2 such that $Nm_{f_2}(F) \in |\delta_2|$. Assume that $cl(\pi_2^*F) \in SingZ_2$. By Proposition (2.1) this is equivalent to $h^0(\tilde{C}, \pi_2^*F) \geq 2$. Then we claim that $h^0(C, F) \geq 2$, so π_2^*F belongs to the locus (23). Indeed, since $\pi_2 : \tilde{C} \rightarrow C$ is a double unramified covering corresponding to $f_2^*(\epsilon) \in Pic^0(C_2)_2$ we have

$$h^0(\tilde{C}, \pi_2^*F) = h^0(C_2, F) + h^0(C_2, f_2^*(\epsilon)(F)) \quad (24)$$

By Lemma (0.1) we conclude that $h^0(C_2, f_2^*(\epsilon)(F)) = 0$. So, $cl(\pi_2^*(F))$ belongs to $Sing(Z_2)$ if and only if $cl(F) \in W_{g-1}^1(C_2)$.

Now, suppose that $F = f_2^*(x) + G$ where $x \in E, G$ is effective and $Nm_{f_2}(G) \in |\delta_2 - 2x|$. Let $t_\epsilon(x)$ be the translation of x by ϵ . Then by Eq. (24) we have

$$h^0(\tilde{C}, \pi_2^*F) = h^0(C_2, f_2^*(x) + G) + h^0(C_2, f_2^*(t_\epsilon(x)) + G) \quad (25)$$

Thus $h^0(\tilde{C}, \pi_2^*F) \geq 2$ and $cl(\pi_2^*F) \in SingZ_2$. The sublocus of $SingZ_2$

$$V = \{cl(\pi_2^*(f_2^*(x) + G)) \mid x \in E, G \geq 0, Nm_{f_2}(G) \in |\delta_2 - 2x|\}$$

is the image of X where X is defined by the pull-back diagram

$$\begin{array}{ccc} X & \longrightarrow & E \\ \downarrow & & \downarrow \beta \\ C_2^{(g-3)} & \xrightarrow{\alpha} & J_2(E) \end{array} \quad (26)$$

and $\alpha(G) = cl(\delta_2 - Nm_{f_2}(G)), \beta(x) = cl(2x)$. The same argument as in Claim 1 of Lemma (5.1) shows that X is irreducible. This implies that V is irreducible as well. Corollary (0.1) and Eq. (25) show that for $F = f_2^*(x) + G$ one has $h^0(\pi_2^*F) = 2$ if and only if G is f_2 -simple. Thus the points (21) form a Zariski open, dense subset of V .

Finally, $dimX = g - 3$ and we claim that the map $X \rightarrow V$ given by $(G, x) \mapsto cl(\pi_2^*(f_2^*(x) + G))$ is of degree 2, hence $dimV = g - 3$. Indeed, let $\sigma_2 : \tilde{C} \rightarrow \tilde{C}$ be the

involution which interchanges the sheets of π_2 . Then for any $\pi_2^*(\xi_2) \in V$ with $h^0(\tilde{C}, \pi_2^*(\xi_2)) = 2$ according to Eq. (25) there are exactly two σ_2 -invariant divisors in $|\pi_2^*(\xi_2)|$ namely

$$\pi_2^*(f_2^*(x) + G), \quad \pi_2^*(f_2^*(t_\epsilon(x)) + G) \quad (27)$$

with x, G as in the lemma. q.e.d.

Let

$$S_2 = \{F \in C_2^{(g-2)} \mid Nm_{f_2}(F) \in |\delta_2|\}$$

One has a surjective map

$$\varphi = cl \circ \pi_2^* : S_2 \longrightarrow Z_2$$

From the proof of Claim 2 of Lemma (5.1) we see that S_2 is irreducible. Moreover $\deg \varphi = 1$ since $h^0(\tilde{C}, L) = 1$ for any sufficiently general $L \in Z_2$. Let us consider the Stein factorization [H]

$$\varphi = \psi \circ \alpha : S_2 \longrightarrow \Gamma \longrightarrow Z_2$$

where ψ is a finite map and α has connected fibers. Let

$$W_1 = \{F \in S_2 \mid h^0(C_2, F) \geq 2\}, \quad W_2 = \{f_2^*A + E \in S_2 \mid A \geq 0, E \geq 0, \deg(A) \geq 2\}$$

One has $\text{codim}_{S_2}(W_1) \geq 1$ and $\text{codim}_{S_2}(W_2) \geq 2$. Let $S_2^0 = S_2 \setminus (W_1 \cup W_2)$ and let $\Gamma^0 = \alpha(S_2^0)$.

Lemma 5.4 *The points of S_2^0, Γ^0 are nonsingular in S_2, Γ respectively, $\text{codim}_\Gamma(\Gamma \setminus \Gamma^0) \geq 2$ and the map $\alpha : S_2^0 \longrightarrow \Gamma^0$ is an isomorphism. Let $n : N \longrightarrow Z_2$ be the normalization of Z_2 . Then there exists a finite map $\beta : N \longrightarrow \Gamma$ such that $n = \psi \circ \beta$. If $N^0 = \beta^{-1}(\Gamma^0)$ then $\text{codim}_N(N \setminus N^0) \geq 2$ and $\beta : N^0 \longrightarrow \Gamma^0$ is an isomorphism*

Proof. The points of S_2^0 are nonsingular by Proposition (0.1). For any $x \in S_2^0$ one has $\#\varphi^{-1}(\varphi(x)) = 2$ as we have shown in the proof of Lemma (5.3). Thus the map $\alpha : S_2^0 \longrightarrow \Gamma^0$ is bijective. The map φ is nondegenerate at the points of S_2^0 . Indeed $\varphi = cl \circ \pi_2^* = \pi_2^* \circ cl$, the map $cl : S_2 \longrightarrow J_{g-1}(C_2)$ is nondegenerate at any $F \in S_2^0$ since $h^0(F) = 1$, and the map $\pi_2^* : J_{g-1}(C_2) \longrightarrow J_{2g-2}(\tilde{C})$ is obviously nondegenerate. We conclude that $\alpha : S_2^0 \longrightarrow \Gamma^0$ is an isomorphism. One has $\text{codim}_\Gamma(\Gamma \setminus \Gamma^0) \geq 2$ since $\text{codim}_{Z_2} \varphi(W_1) \geq 2$ by (22). The rest of the lemma is clear by the universal property of the normalization. q.e.d.

Reconstruction of (C, η) in the bi-elliptic case, $g \geq 4$.

In the beginning of this section we have seen how to reconstruct up to isomorphism E and the covering $f_2 : C_2 \longrightarrow E$. Let $T_i \in O_i \subset |2\Xi|$, $i = 1, 2$. We have proved above that T_i are irreducible and just one of T_i has a singular locus of codimension 1. Reordering, if necessary, as in Lemma (5.1) we can assume that this divisor is T_2 . We can identify T_2 and Z_2 by translation. Let $n : N \longrightarrow T_2$ be the normalization of T_2 . Let $R = n^{-1}(V)$. The Zariski open subset $R^0 = R \cap N^0$ is dense in R since $\text{codim}_N(N \setminus N^0) \geq 2$. By the irreducibility of X in (26) one concludes that $\alpha^{-1} \circ \beta(R^0)$ and R are irreducible as well. We have an isomorphism $f^* : |\delta_2| \longrightarrow |K_C \otimes \eta|$. Consider the Gauss map $G : Z_2^{ns} \longrightarrow |K_C \otimes \eta|$. Then for every $F \in S_2$ with $h^0(\tilde{C}, \pi_2^* F) = 1$ one has

$$G(\varphi(F)) = f^*(Nm_{f_2}(F))$$

Shrinking N_0 from Lemma (5.4) we can assume that the following properties hold

- $\text{codim}_N(N \setminus N_0) \geq 2$.
- The composition $C \circ n$ can be extended to a regular map on N_0 .
- Every point of $\alpha^{-1} \circ \beta(R^0)$ has the form $f_2^*(x) + A$ where x is not a branch point of f_2 , A is reduced and f_2 -simple, and $x \notin \text{Supp}(A)$.

Now, we define a rational map

$$\gamma : R \longrightarrow E$$

as follows. For every $L \in R_0$ the hyperplane $G \circ n(L)$ belongs to the unique irreducible component of degree > 1 of the branch locus of the Gauss map $G : T \longrightarrow \mathbf{P}(T_0 P)^*$, namely E^* . By the conditions above this hyperplane is tangent to a unique point of E . We denote this point by $\gamma(L)$. Now, let $L = \beta^{-1} \circ \alpha(f_2^*(x) + A) \in R_0$. Then $n^{-1}(n(L)) = \{L, L'\}$ where

$$L' = \beta^{-1} \circ \alpha(f_2^*(t_\epsilon(x)) + A) \quad (28)$$

according to (27). This shows that the map $n : R \longrightarrow V \subset \text{Sing}(T_2)$ is of degree 2. By (28) the corresponding involution $\tau^* : \mathbf{C}(R) \longrightarrow \mathbf{C}(R)$ of the field of rational functions on R transforms $\gamma^* \mathbf{C}(E)$ into itself and $\tau^* : \gamma^* \mathbf{C}(E) \longrightarrow \gamma^* \mathbf{C}(E)$ is induced by the translation map $t_\epsilon : E \longrightarrow E$. This gives the reconstruction of $\epsilon \in J(E)_2$ and completes the reconstruction of (C, η) from the extended Prym data.

6 The case $g = 3$

Let $a \in \tilde{C}$. We define the Abel-Prym map $\phi : \tilde{C} \longrightarrow P(\tilde{C}, \sigma)$ by

$$\phi(x) = cl(x - a - \sigma(x - a))$$

Lemma 6.1 *Suppose $g \geq 2$. The following alternative takes place*

(i) *ϕ maps $\tilde{C}$ isomorphically onto its image $\phi(\tilde{C})$.*

(ii) *The map $\phi : \tilde{C} \longrightarrow \phi(\tilde{C})$ has degree 2.*

The second case occurs if and only if

(*) *C is hyperelliptic and $\eta \simeq \mathcal{O}_C(p_1 - p_2)$ for some $p_1, p_2 \in C$.*

Here $\phi(\tilde{C}) \simeq C_2$ and $\phi = \pi_2$ (see Fig.1) via this isomorphism.

Proof. Suppose $\phi(x) = \phi(y)$ for some $x \neq y$. Then $x + \sigma y \equiv y + \sigma x$, thus $\tilde{C}$ is hyperelliptic. It has a unique g_2^1 , so $\sigma(g_2^1) = g_2^1$. Let σ_1 be the hyperelliptic involution of $\tilde{C}$. Then $\sigma \neq \sigma_1$ and we claim that σ and σ_1 commute. Indeed, for any $z \in \tilde{C}$

$$\sigma z + \sigma_1(\sigma z) \in g_2^1, \quad \sigma(z + \sigma_1 z) \in g_2^1$$

Thus $\sigma\sigma_1 z = \sigma_1\sigma z$. Let $\sigma_2 = \sigma\sigma_1$. Let $C_1 = \tilde{C}/\sigma_1$ and let $\bar{\sigma} : C_1 \longrightarrow C_1$ be the involution induced by σ . Then $\bar{\sigma}$ has two fixed points since $C_1 \simeq \mathbf{P}^1$. Thus $\pi : \tilde{C} \longrightarrow C$ fits into the commutative diagram of Fig.1 and condition (*) holds. If ϕ were degenerate at some point $x \in \tilde{C}$, then $\pi(x)$ would be a base point of $|K_C \otimes \eta|$, thus condition (*) holds according to Lemma (3.1). Conversely, suppose condition (*) holds. Then by the argument above $\phi(x) = \phi(y)$ and $x \neq y$ if and only if y belongs to the divisor $\sigma(x + \sigma_1 x)$. Thus $y = \sigma_2(x)$. q.e.d.

Further we suppose that $g = 3$. Let $T_i \in \mathcal{O}_i \subset |2\Xi|, i = 1, 2$. The divisors T_i are reduced, connected curves according to Proposition (2.1). Let $S = Nm^{-1}(|K_C \otimes \eta|) \subset \tilde{C}^{(4)}$ and let $Z = Nm^{-1}(K_C \otimes \eta) \cap W_4(\tilde{C}) \subset J_4(\tilde{C})$. Both S and Z break into two disjoint, connected components $S = S_1 \cup S_2, Z = Z_1 \cup Z_2$. We enumerate so that $cl(S_i) = Z_i$ and T_i is translation of $Z_i, i = 1, 2$.

Lemma 6.2 *The curves T_1, T_2 are both singular if and only if condition (*) of Lemma (6.1) holds. If only one of T_i is singular then the nonsingular one is a translation of $\phi(\tilde{C})$.*

Proof. If condition (*) holds, then both T_1 and T_2 are reducible and hence singular by Corollary (4.1). Since T_i is a translation of $Z_i, i = 1, 2$ we can work with Z_i . Suppose that condition (*) does not hold and $Z' \in \{Z_1, Z_2\}$ is a singular curve with a singular point L . Let $\{Z', Z''\} = \{Z_1, Z_2\}$. Then

$$X = \{L + x - \sigma x \mid x \in \tilde{C}\} \subset Z''$$

Clearly X is a translation of $\phi(\tilde{C})$. According to Lemma (6.1) X is isomorphic to $\tilde{C}$ and X is algebraically equivalent to 2Ξ [Ma]. Thus $X = Z''$. q.e.d.

Reconstruction of (C, η) in the case $g = 3$.

Case 1. *Both T_1, T_2 are singular.*

According to Lemma (6.2) we are in the situation of Section (4) where a procedure for the reconstruction of (C, η) was described.

Case 2. *Just one of the curves $T' \in \{T_1, T_2\}$ is singular.*

We take the other curve T'' . It is isomorphic to $\tilde{C}$ according to Lemmas (6.2) and (6.1). The involution $-id_P : T'' \rightarrow T''$ coincides with σ via this isomorphism. We thus reconstruct $(\tilde{C}, \sigma)$.

Case 3. *T_1 and T_2 are nonsingular.*

Consider the involutions $\sigma_i : T_i \rightarrow T_i$ induced by $-id_P$, let $C_i = T_i/\sigma_i$ and let $\pi_i : T_i \rightarrow C_i$ be the factor maps, $i = 1, 2$. By Proposition (2.1) the map $cl : S_i \rightarrow Z_i$ is bijective. Thus S_i are nonsingular since Z_i are nonsingular. Using Proposition (0.1) one checks that the nonsingularity of S_i implies that $\sigma^{(4)} : S_i \rightarrow S_i$ is without fixed points, thus $\sigma_i : T_i \rightarrow T_i$ is without fixed points as well, $i = 1, 2$. Consider the Gauss maps $G_i : T_i \rightarrow \mathbf{P}(T_0P)^* = \mathbf{P}^1$. By Eq. (9) one shows that $G_i = f_i \circ \pi_i$ where $f_i : C_i \rightarrow \mathbf{P}^1$ are maps of degree 4 and concludes that the maps $G_i : T_i \rightarrow \mathbf{P}^1$ are obtained from $\varphi_{K_C \otimes \eta} \circ \pi : \tilde{C} \rightarrow \mathbf{P}^1$ by the tetragonal construction of Donagi [Do1],[Do2]. Now, take the pair (T_1, σ_1) and apply the tetragonal construction to $G_1 : T_1 \rightarrow \mathbf{P}^1$ (ibid.). One obtains two 8-sheeted coverings $g_i : X_i \rightarrow \mathbf{P}^1$ with involutions $\tau_i : X_i \rightarrow X_i$ such that $g_i \circ \tau_i = g_i$. For one of them, e.g.

X_2 , there is an isomorphism of the coverings

$$\begin{array}{ccc} X_2 & \xrightarrow{\psi_2} & T_2 \\ g_2 \downarrow & & \downarrow G_2 \\ \mathbf{P}^1 & = & \mathbf{P}^1 \end{array}$$

such that $\psi_2 \circ \tau_2 = \sigma_2 \circ \psi_2$. Then the remaining pair (X_1, τ_1) is isomorphic to $(\tilde{C}, \sigma)$.

References

- [A] A. Andreotti, *On a theorem of Torelli*, Amer. J. Math. **80** (1958), 801-828.
- [ACGH] E. Arbarello, M. Cornalba, P. A. Griffiths & J. Harris, *Geometry of algebraic curves*, Vol. I, Springer-Verlag, Berlin, 1985.
- [B] A. Beauville, *Fibrés de rang 2 sur une courbe, fibré déterminant et fonctions thêta*, Bull. Soc. Math. France **116** (1988), 431-448.
- [C] C. H. Clemens, *A scrapbook of complex curve theory*, Plenum Press, New York, 1980.
- [Da] S. G. Dalaljan, *The Prym variety of two-sheeted covering of a hyperelliptic curve*, Uspehi Mat. Nauk. **29** (1974) no. 6, (180), 165-166 (in Russian).
- [De1] O. Debarre, *Sur les variétés de Prym des courbes tétraogales*, Ann. Sci. E.N.S. **21** (1988), 545-559.
- [De2] O. Debarre, *Le théorème de Torelli pour les intersections de trois quadriques*, Invent. Math. **95** (1989), 507-528.
- [Do1] R. Donagi, *The tetragonal construction*, Bull. Amer. Math. Soc. (N.S.) **4** (1981), 181-185.
- [Do2] R. Donagi, *The fibers of the Prym map*, Contemp. Math. **136** (1992), 55-125.
- [F] J. Fay, *Theta functions on Riemann surfaces*, Lecture notes in Mathematics Vol.332, Springer-Verlag, Berlin, 1973.

- [F-S] R. Friedman, R. Smith, *The generic Torelli theorem for the Prym map*, Invent. Math. **75** (1984), 85-104.
- [H] R. Hartshorne, *Algebraic geometry*, Springer-Verlag, Berlin, 1977.
- [H-K] A. Hefez, S. L. Kleiman, *Notes on the duality of projective varieties*, Progress in Math. Vol. 60: Geometry Today, Birkhäuser, Boston, 1985, pp.143-193.
- [K] V. Kanev, *The global Torelli theorem for Prym varieties at a generic point*, Math USSR Izvestija **20** (1983), 235-258.
- [Ma] L. Masiewicki, *Universal properties of Prym varieties with application to algebraic curves of genus five*, Trans. Amer. Math. Soc. **222** (1976), 221-240.
- [M1] D. Mumford, *Theta characteristics of an algebraic curve*, Ann. Sci. E.N.S. **4** (1971), 181-192.
- [M2] D. Mumford, *Prym varieties I*, in Contributions to Analysis (Eds. L. V. Ahlfors et al.), Academic Press, New York, 1974, pp.325-350.
- [M3] D. Mumford, *Tata lectures on theta* Vol. I, Progress in Math. Vol. 28, Birkhäuser, Boston, 1983.
- [N] J. C. Naranjo, *Prym varieties of bi-elliptic curves*, J. reine angew. Math. **B. 424** (1992), 47-106.
- [T] A. N. Tjurin, *The geometry of the Poincaré theta divisor of a Prym variety*, Math USSR Izvestija **39** (1975), 951-986.
- [V] A. Verra, *The Prym map has degree two on plane sextics*, Preprint, 1992, 26 p.
- [W] G. E. Welters, *Abel-Jacobi isogenies for certain types of Fano threefolds*, Thesis, Mathematisch Centrum, Amsterdam, 1981.

INSTITUTE OF MATHEMATICS, BULGARIAN ACADEMY OF SCIENCES,
 Acad. G. Bonchev Str. bl. 8, SOFIA 1113 BULGARIA